

A Golf Course Bibliography

By

Steve Klaine

Golf is a multimillion dollar business that requires the use of enormous amounts of pesticides and fertilizers to maintain the well manicured course that is expected by those paying to play. Each course, averaging between 124 and 180 acres (Beard, 1982; Watson, 1990) is managed by a superintendent that is educated in the art of turfgrass maintenance. The turfgrass industry includes not only golf courses but other sports playing fields and the home lawn. Of this huge industry, golf only represents approximately 6% of the total money involved (Cockerman, 1985; Watson, 1990). Different regions of the country require specific management strategies due to differences in climate, geology, and hydrology. Therefore, superintendents have their own strategies for their course that was developed from experience and various other sources such as the scientific literature.

Public concerns for the environment and increasingly stringent water quality restrictions have led to the idea of best management practices (BMPs) for golf courses. Krivak (1978) conceptualized BMPs for agricultural crops. The BMP concept consist of five basic goals: 1) decrease the offsite transport of pesticides and nutrients, 2) control the application of these chemicals, 3) decrease the total chemical loads, 4) use both biological and mechanical soil and water conservation plans (SWCPs), and 5) educate both the managers and the public about the relationship between environmental issues and golf course management (Balogh and Walker, 1992).

However, research published to date lacks the ability to provide BMPs to golf course superintendents. There are only a few attempts at such a task (Leslie and Metcalf, 1989; Walker, 1990; USDA-ARS, 1988a; USDA-ARS, 1988b). The project at the Ocean Course on Kiawah Island is an effort to bring together many of facets of environmental research into a single holistic approach to providing plausible management strategies. Another study currently being undertaken at Clemson University is a comprehensive survey of all golf course managers in South Carolina. Compilation and comparison of each of the various strategies will hopefully provide a base to build best management practices for superintendents.

Balogh and Walker (1992) have outlined the potential detriments of construction and management of golf courses. Probably the most important concern of the present is the potential for contamination of surface and groundwaters with sediment, nutrients, and pesticides (Cooper, 1987; Grant, 1987; Klien, 1990; EPA, 1986; EPA 1988; Keeney, 1986; Petrovic, 1990; Pratt, 1985; Pye *et al.*, 1983). Continuous use of pesticides in turfgrass management strategies may also lead to an increased resistance by the target organisms (Potter and Braman, 1991). Obviously, there is the concern of possible negative impacts on nontarget organisms by chemical management strategies (Kendall *et al.*, 1992; Stone and Gardoni, 1985a; Stone and Gardoni, 1985b; Stone, 1979). The U.S. is not the only place where public awareness has brought this to governmental agencies attention. Japan has begun setting standards for maximum allowable pesticide residues found in golf course effluent (Tsuda *et al.*, 1992). In order to maintain the lush green carpet of turfgrass, an extremely

large quantity of water is used. The push for water conservation has resulted in increased utilization of secondary treated sewage effluent for irrigation (Roberts, 1989; Payne, 1987). Effects of golf courses on the precious wetlands of the U.S. have also been of public concern. Developers and managers have also realized their importance (Dye, 1989; Peacock, 1990; Salvesen, 1990). With the increasing cost of "after the fact" mitigation, better management practices that prevent some of these problems are being evoked. Slowly, scientific research on environmental impacts of golf courses are being published (Leslie and Metcalf, 1989; Petrovic, 1990; Walker, 1990). However, much of the past and present research presents only a small portion of the whole picture. There is a great need for more comprehensive studies and literature reviews (Balogh and Walker, 1990; Beard, 1982; Beard, 1973; Madison, 1982). Computer expert systems for planning turf management such as TURFPLAN appear to work well for low maintenance applications, but fall short of designs of human experts for high maintenance turfgrass such as golf courses (Liu et al., 1991).

In the battle to establish management strategies for golf courses, there have been two basic approaches. Some claim that the total acreage maintained (Smolen, 1984) is a better indicator of possible risks as opposed to the obvious quantitative total volume of chemicals applied. This partly stems from the fact that the total annual use and number of pesticides or fertilizers applied varies among courses and within courses (Beard, 1982). There are three basic management practices concerned with golf courses that raise environmental issues. Pest control involves the use of a wide

range of chemicals including insecticides, herbicides, and fungicides. Plant growth regulators (PGRs) may also fall into this category because of their basic chemical makeup. An alternative to the use of pesticides that has some promise is that of biological pest controls (Myers *et al.*, 1992). The cultural practices of fertilization and irrigation represent the remaining two that gain public attention. Both have been shown to affect pest populations and subsequent damage (Beard, 1973). Nitrate (NO_3) originating from cess pools, septic tanks, animal and human wastes, and fertilizers (Keeney, 1986) is one of the most widespread groundwater contaminants (Pye *et al.*, 1983). Finally, irrigation and drainage practices directly affect pesticide and nutrient applications in turfgrass management (Biran, 1981; Colbaugh, 1985). One can now see that all management decisions are interrelated and codependent.

Probably foremost in the mind of a superintendent when planning pesticide management is the safety of the workers and golfers. A few studies have made attempts at finding safe levels of dislodgeable residues and the time required post-application to achieve them (Goh *et al.*, 1986; Harris and Soloman, 1992). The results indicate that for most pesticides used, there is little risk involved with reentry to a treated site. A method for quantifying airborne loss of pesticides was described by Jenkins *et al.* (1991). The only practical method for preventing serious damage of turf by insects is the use of insecticides (Potter and Braman; 1991). Insecticides may have indirect effects that counteract or cause a more serious problem. They have been shown to adversely affect earthworms causing thatch buildup (Randell *et al.*, 1972).

Populations of predators and parasitoids in the soil may also be diminished causing some secondary outbreak (Cockfield, 1983). Conservation of these natural enemies should factor into pesticide selection. Certain commonly used herbicides also pose the threat of causing phytotoxicity and decreased turf quality. The timing of applications to sensitive turf species such as creeping bentgrass is very important (Shim and Johnson, 1992). Frequent use of select fungicides may enhance some nontarget diseases, while the nontarget benefits appear to be turf species dependent (Dernoenden and McIntosh, 1991).

The use of plant growth regulators for golf course applications are limited due to the potential for turf damage and inconsistent results (Christians, 1985). Some PGRs have been found effective if used in areas that are hazardous for trimming or mower operation (Fry, 1991). Fry also reported that glyphosate (0.6 kg/ha) caused unacceptable phytotoxicity.

Johnson (1992) reported variations in effects, from slight to moderate turf damage by two PGRs. Recovery was complete by 10 weeks. This would appear unacceptable for the golf industry that is forced to maintain turf at the highest quality possible. Hence, more research is needed to pronounce sentence on PGRs. Outside of these pure turf management concerns, the superintendent must also be environmentally sensitive and be concerned about pesticide runoff. Binding to thatch, which increases retention time for degradation processes, is reported to diminish the amounts of pesticides in runoff to safe levels in receiving waters (Miles *et al.*, 1992; Potter and Braman, 1991; Watschke, 1990). Enhancement of natural phenomena is a management technique that is receiving more attention.

Cranshaw and Zimmerman (1989) reported effective results when using nematodes to control turfgrass scarabs. They also go on to point out problems associated with this practice such as: availability, storage, cost, handling, and reliability. Dollar spot, a common disease found on golf courses (Smiley, 1983) was effectively suppressed with applications of compost (Nelson and Craft, 1992). This research was based on the fact that composted substrates hold disease-suppressive properties due to their microbial content (Hoitink and Fahy, 1986). Again, more field research is needed to evaluate the usefulness of biological control methods for use on golf courses.

Fertilizer management poses some concerns similar to those associated with pesticides. Petrovic (1990) summarizes and reviews the literature dealing with nitrogenous fertilizer usage. Nitrogen and potassium fertilization of turfgrass has been linked to resistance to stress conditions brought on by disease, drought, or human foot traffic (Beard, 1973; Cook *et al.*, 1983; Markland, 1969). The distribution of nitrogenous fertilizers is normally studied as a series of components rather than a complete system. Even though conclusions are limited to a certain cultural and geological situation, Starr and DeRoo (1981) attempted to study all of the components. Atmospheric loss of nitrogen may occur via NH_3 vaporization or denitrification. Ammonia volatilization can be decreased by irrigation (Bowman *et al.*, 1987), decreased thatch content, and the use of time-release nitrogen pellets (Nelson *et al.*, 1980). The process of denitrification and its effects are limited in the literature (Mancino *et al.*, 1988). Controlled-release or time-release fertilizers are used commonly on

turfgrass (Turgeon, 1985). A study of reactive layer coated (RLC) nitrogen (Peacock and DiPaola, 1992) yielded results indicating that their effectiveness depends upon the thickness of the reactive layer coating. A similar study involving controlled-release potassium (Snyder and Cisar, 1992) looked at the coating material's effect on potassium release. Sulfur-coated (SC) released potassium too rapidly, while the converse was true for resin-coated (RC) potassium. All other sources of potassium tested had favorable characteristics. As with pesticides, leaching and runoff are also a concern. Leaching is highly influenced by: soil texture, nutrient source, rate and timing of application, and irrigation/rainfall (Petrovic, 1990). It is thought that because a turfgrass ecosystem results in soils with high infiltration capacity, runoff is seldom a problem (Gross *et al.*, 1990; Gross *et al.*, 1991).

Obviously a big factor in the previous discussions, water, via rainfall or irrigation, must be seriously considered when formulating a management strategy. Organophosphates and carbamates applied in a granule form require immediate irrigation so as to reach the target organism as well as decrease the chances of nontarget exposure (Potter and Braman, 1991). Just as adequate fertilization enhances turfgrass quality and resistance, so does maintaining the appropriate soil moisture (Couch and Bloom, 1960; Sugimoto *et al.*, 1990). A chronic study (3.3 yrs) monitoring the soil quality of turfgrass irrigated with secondary treated effluent was conducted by Mancino and Pepper (1992). Soil salinity, pH, Fe, K, Na, and P levels all increased. Addition of calcium sulfate or sulfur can be used to bring the sodium

levels back to normal. The excess phosphorous could be accounted for by decreasing P added in the form of fertilizers. No effect was recorded on the levels of total organic carbon (TOC), total nitrogen (TN), or total aerobic bacteria. So it seems that irrigation with effluent may require additional management for some things while eliminating others. Nus and Sandburg (1991) point out that irrigation with effluent must be closely monitored. Phytotoxic chemicals such as herbicides (eg. atrazine) may be present at levels to cause chronic problems.

In conclusion, much work still has to be done in order to present golf course superintendents with a "cookbook" approach to pest management. An interdisciplinary team consisting of superintendents, agronomist, environmental toxicologist, and chemist are needed to review past and current strategies. From this intensive cooperation, future research and strategies may be formed.

Selected References

- Balogh, J.C. and W.J. Walker. 1992. Golf Course Management and Construction: Environmental Issues. Lewis Publishers, Ann Arbor, MI, 951 pp.
- Beard, J.B.. 1982. Turf Management for Golf Courses. Macmillan Publishing Co., New York, 642 pp.
- Beard, J.B.. 1973. Turfgrass Science and Culture. Prentice-Hall, Inc., Englewood Cliffs, NJ, 658 pp.
- Bezella, D.A. 1975. Let's manage an irrigation pond on a golf course and control aquatic weeds. *Park Maintenance* 28(4): 10-13.
- Biran, I., B. Bravdo, I. Bushkin-Harav, and E. Rawitz. 1981. Water consumption and growth rate of 11 turfgrasses as affected by mowing height, irrigation frequency, and soil moisture. *Agronomy Journal* 73(1): 85-90.
- Bowman, D.C., J.L. Paul, W.B. Davis, and S.H. Nelson. 1987. Reducing ammonia volatilization from Kentucky bluegrass turf by irrigation. *Horticulture Science* 22: 84-87.
- Cockerham, S.T. and V.A. Gibeault. 1985. The size, scope, and importance of the turfgrass industry. In: Gibeault, V.A. and S.T. Cockerham (Eds.). Turfgrass Water Conservation. University of California, Riverside, Division of Agriculture and Natural Resources, pp. 7-12.
- Cockfield, S.D. and D.A. Potter. 1983. Short-term effects of insecticidal applications of predaceous arthropods and oribatid mites in Kentucky bluegrass turf. *Environmental Entomology* 12: 1260-1264.
- Cohen, S.H., S. Nickerson, R. Maxey, A. Dupuy, and J.A. Senita. 1990. A ground water monitoring study for pesticides and nitrates associated with golf courses on Cape Cod. *Ground Water Monitoring Review* 10(1): 160-173.

- Cook, R.J. and K.F. Baker. 1983. The Nature and Practice of Biological Control of Plant Pathogens. American Phytopathological Society, St. Paul, MN.
- Cooper, R.J.. 1987. Turfgrass pesticide risk in perspective. *Golf Course Management* 55(11): 50-60.
- Cranshaw, W.S. and R.J. Zimmerman. 1989. Biological, mechanical, and chemical control of turfgrass-infesting scarabs in Colorado. *Southwestern Entomologist* 14(4): 351-355.
- Christians, N.E. 1985. Response of Kentucky bluegrass to four growth retardants. *Journal of the American Society for Horticulture Science* 110: 765-769.
- Dernoeden, P.H. and M.S. McIntosh. 1991. Seasonal quality responses of perennial ryegrass as influenced by fungicides. *Hortscience* 26(9): 1181-1183.
- Dye, P.. 1989. Golf course architects adapt to environmental challenges. *SportsTURF* 5(5): 29.
- Edmonson, J. 1987. Hazards of the game. *Audubon* 89(6): 24-27.
- Environmental Protection Agency. 1986. Pesticides in Ground Water: Background Document. EPA-WH550G. Office of Ground-Water, U.S. Environmental Protection Agency, Washington, D.C., 72 pp.
- Environmental Protection Agency. 1988. Protecting Ground Water: Pesticides and Agricultural Practices. EPA-440/6-88-001. U.S. Environmental Protection Agency, Office of Ground-Water Protection, Washington, D.C., 53 pp.
- Fry, J.D. 1991. Centipedegrass response to plant growth regulators. *Hortscience* 26(1): 40-42.
- Giblin Davis, R.M., J.L. Cisar, F.G. Bilz, and K.E. Williams. 1991. Management practices affecting phytoparasitic nematodes in Tifgreen Bermudagrass. *Nematropica* 21(1): 59-69.

- Goh, K.S., S. Edmiston, K.T. Maddy, D.D. Meinders, and S. Margetich. 1986. Dissipation of dislodgeable foliar residue of Chlorpyrifos and Dichlorvos on turf. *Bulletin of Environmental Contamination and Toxicology* 37 (1): 27-33.
- Grant, Z.. 1987. One aspect of government relations: groundwater issues in pesticide regulation. *Golf Course Management* 55(6): 14-20.
- Gross, C.M., J.S. Angle, R.L. Hill, and M.S. Welterlen. 1991. Runoff and sediment losses from tall fescue under simulated rainfall. *Journal of Environmental Quality* 20(3): 604-608.
- Gross, C.M. 1990. Nutrient and sediment losses from turfgrass. *Journal of Environmental Quality* 19(4): 663-669.
- Harris, S.A. and K.R. Solomon. 1992. Human exposure to 2,4-D following controlled activities on recently sprayed turf. *Journal of Environmental Science and Health (Part B): pesticides, food contaminants, and agricultural wastes*. 27(1): 9-22.
- Hoitink, H.A.J. and P.C. Fahy. 1986. Basis for the control of soilborne plant pathogens with composts. *Annual Review of Phytopathology* 24: 93-114.
- Horsley, S.W. and J.A. Moser. 1990. Monitoring ground water for pesticides at a golf course: a case study on Cape Cod, Massachusetts. *Ground Water Monitoring Review* 10(1): 101-108.
- Jenkins, J.J., R.J. Cooper, and A.S. Curtis. 1991. Field chamber technique for measuring Pendimethalin airborne loss from turfgrass. *Bulletin of Environmental Contamination and Toxicology* 47(4): 594-601.
- Johnson, B.J. 1992. Response of Tifway Bermudagrass to rate and frequency of Flurprimidol and Paclobutrazol application. *Hortscience* 27(3): 230-233.

- Keeney, D.R.. 1986. Sources of nitrate to groundwater. CRC Critical Reviews of Environmental Control 16: 257-304.
- Kendall, R.J., L.W. Brewer, R.R. Hitchcock, and J.R. Mayer. 1992. American wigeon mortality associated with turf application of Diazinon AG500. Journal of Wildlife Diseases 28(2): 263-267.
- Klein, R.D.. 1990. Protecting the Aquatic Environment from the Effects of Golf Courses. Community & Environmental Defense Associates, Maryland Line, MD, p. 59.
- Krivak, J.A.. 1978. Best management practices to control nonpoint-source pollution from agriculture. Journal of Soil and Water Conservation 33(4): 161-166.
- Leslie, A.R. and R.L. Metcalf (Eds.). 1989. Integrated Pest Management for Turfgrass and Ornamentals. Office of Pesticide Programs, 1989-626-030. U.S. Environmental Protection Agency, Washington, D.C., 337 pp.
- Liu, H., T.W. Fermanian, and S.G. Carmer. 1991. Expert system for planning the establishment of turfs. Agronomy Journal 83(1): 140-143.
- Madison, J.H.. 1982. Principles of Turfgrass Culture. Robert E. Krieger Publishing Co., Melbourne, FL, 431 pp.
- Maschek, M.E. 1989. SCADA system employed in water reuse project. Water Engineering and Management 136(8): 29-32.
- Mancino, C.F. and I.L. Pepper. 1992. Irrigation of turfgrass with secondary sewage effluent: soil quality. Agronomy Journal 84(4): 650-654.
- Mancino, C.F., W.A. Torello, and D.J. Wehner. 1988. Denitrification losses from Kentucky bluegrass sod. Agronomy Journal 80: 148-153.
- Markland, F.E., E.C. Roberts, and L.R. Frederick. 1969. Influence of nitrogen fertilizers on Washington creeping bentgrass, *Agrostis palustris* Huds. II. Incidence of dollar spot, *Sclerotinia homoeocarpa*, infection. Agronomy Journal 61: 701-705.

- McCarty, L.B., L.T. Lucas, and J.M. Dipaola. 1992. Spring dead spot occurrence in bermudagrass following fungicide and nutrient applications. *Hortscience* 27(10): 1092-1093.
- Miles, C.J., G. Leong, and S. Dollar. 1992. Pesticides in marine sediments associated with golf course runoff. *Bulletin of Environmental Contamination and Toxicology* 49(2): 179-185.
- Mujeriego, R. and L. Sala. 1991. Golf course irrigation with reclaimed wastewater. *Water Science and Technology* 24(9): 161-172.
- Mus, J.L. and M.A. Sandburg. 1991. Creeping Bentgrass damaged by low levels of Atrazine in irrigation water. *Hortscience* 26(4): 393-394.
- Myers, R.F., R.E. Wagner, and P.M. Halisky. 1992. Relationship between cultural factors and nematodes on Merion Kentucky Bluegrass. *Journal of Nematology* 24(1): 205-211.
- Nelson, E.B. and C.M. Craft. 1992. Suppression of dollar spot on Creeping Bentgrass and Annual Bluegrass turf with compost-amended topdressings. *Plant Disease* 76(9): 954-958.
- Nelson, K.E., A.J. Turgeon, and J.R. Street. 1980. Thatch influence on mobility and transformation of nitrogen carriers applied to turf. *Agronomy Journal* 72: 487-492.
- Payne, R.A.. 1987. Resource conservation: the golf course role. *Golf Course Management* 55(10): 50-58.
- Peacock, C.H. and J.M. Dipaola. 1992. Bermudagrass response to reactive layer coated fertilizers. *Agronomy Journal* 84(6): 946-950.
- Peacock, C.H., A.H. Bruneau, and S.P. Spak. 1990. Protecting a valuable resource: preservation of wetlands from a technical perspective. *Golf Course Management* 58(11): 6-7.

- Petrovic, A.M. 1990. The fate of nitrogenous fertilizers applied to turfgrass. *Journal of Environmental Quality* 19(1): 1-15.
- Potter, D.A. and S.K. Braman. 1991. Ecology and management of turfgrass insects. *Annual Review of Entomology* 36: 383-406.
- Potter, D.A., M.C. Buxton, C.T. Redmond, C.G. Patterson, and A.J. Powell. 1990. Toxicity of pesticides to earthworms (*Oligochaeta*, *Lumbricidae*) and effect on thatch degradation in Kentucky Bluegrass turf. *Journal of Economic Entomology* 83(6): 2362-2369.
- Pratt, P.F. 1985. Agriculture and Groundwater Quality. Council for Agricultural Science and Technology. CAST Report No. 103, 62 pp.
- Pye, V.I., R. Patrick, and J. Quarles. 1983. Groundwater Contamination in the United States. University of Pennsylvania Press, Philadelphia, 315 pp.
- Randall, R.J., J.D. Butler, and T.D. Hughes. 1972. The effect of pesticides on thatch accumulation and earthworm populations in Kentucky bluegrass turf. *Hortscience* 7: 64-65.
- Roberts, E.C. and B.C. Roberts. 1989. Lawn and Sports Turf Benefits. The Lawn Institute, Pleasant Hill, TN, 31 pp.
- Robinson, C.E. 1971. Golf courses. . . cornerstones of success-ful park operations. *Parks and Recreation* 6(10): 50-54.
- Salvesen, D. 1990. Wetlands: Mitigating and Regulating Development Impacts. The Urban Land Institute, Washington, D.C., 117 pp.
- Sang, R.S. and B.J. Johnson. 1992. Response of Creeping Bentgrass to spring-applied herbicides. *Hortscience* 27(3): 237-239.

- Schreiner, W. 1989. Wetland preservation: par for the course. AWRA Wetlands: Concerns and Successes Symposia (Sept. 17-22). pp. 335-344.
- Shirmohammadi, A., W.L. Magette, and L.L. Shoemaker. 1991. Reduction of nitrate loading to ground water. Ground Water Monitoring Review 11(1): 112-119.
- Shultz, C.D., W.L. Yaeger, and A. Bevenue. 1975. Pesticides in fish from a Hawaiian canal. Bulletin of Environmental Contamination and Toxicology 13(2): 206-209.
- Smiley, R.W.. 1983. Compendium of Turfgrass Diseases. American Phytopathological Society, St. Paul, MN.
- Smolen, M.D., F.J. Humenik, J. Spooner, S.A. Dressing, and R.P. Maas. 1984. Best management Practices for Agricultural Nonpoint Source Control IV Pesticides. USDA Cooperative Agreement 12-05-300-472, EPA Interagency Agreement AD-12-F-0-037-0. North Carolina Agriculture Extension Service, 87 pp.
- Starr, J.L. and H.C. DeRoo. 1981. The fate of nitrogen applied to turfgrass. Crop Science 21: 531-536.
- Stone, W.B. and P.B. Gradoni. 1985. Recent poisonings of wild birds by Diazinon and Carbofuran. Northeastern Environmental Science 4(3-4): 160-165.
- Stone, W.B. and P.B. Gradoni. 1985. Wildlife mortality related to the use of the pesticide Diazinon. Northeastern Environmental Science 4(1): 30-39.
- Stone, W.B. 1979. Poisoning of wild birds by organophosphate and carbamate pesticides. New York Fish and Game Journal 26(1): 37-48.
- Snyder, G.H. and J.L. Cisar. 1992. Controlled-release potassium fertilizers for turfgrass. Journal of the American Society for Horticulture Science. 117(3): 411-414.

- Sugimoto, E.E., H.A.J. Hoitink, and O.H. Tuovinen. 1990. Oligotrophic pseudomonads in the rhizosphere: suppressiveness to *Pythium* damping-off of cucumber seedlings (*Cucumis sativus* L.). *Biol. Fert. Soils* 9: 231-234.
- Tsuda, T., S. Aoki, M. Kojima, and T. Fujita. 1992. Accumulation and excretion of pesticides used in golf courses by carp (*Cyprinus carpio*) and willow shiner (*Gnathopogon caerulescens*). *Comparative Biochemistry and Physiology C-Comparative Pharmacology and Toxicology* 101(1): 63-66.
- Turgeon, A.J. 1985. Turfgrass Management (Revised ed.). Reston Publishing Co., Reston, VA.
- USDA Agricultural Research Service. 1988a. ARS Strategic Groundwater Plan. 1. Pesticides. Washington, D.C.
- USDA Agricultural Research Service. 1988b. ARS Strategic Groundwater Plan. 1. Nitrates. Washington, D.C.
- Walker, W.J., J. Balogh, R.M. Tietge, and S.R. Murphy. 1990. Environmental Issues Related to Golf Course Construction and Management. United States Golf Association, Green Section, Far Hills, NJ, 378 pp.
- Watschke, T.L.. 1990. The environmental fate of pesticides. *Golf Course Management* 58: 18-24.
- Watson, J.R., Jr., H.E. Kaerwer, and D.P. Martin. 1990. The turfgrass industry. In: Turfgrass Science. ASA-CSSA Monogr. 14 (Revised). American Society of Agronomist, Crop Science Society of America, Madison, WI, 106 pp.
- Zimmerman, R.J. and W.S. Cranshaw. 1990. Solutions of pesticides used in turfgrass maintenance. *Journal Economic Entomology* 83(1): 97-100.